

FACT SHEET | SEPTEMBER 2023

Endoscopy and type 2 diabetes

RSS Diabetes Service

An endoscopy involves fasting, changes in your diet, physical activity levels, diabetes medications and may cause stress, anxiety and discomfort. These factors can also disrupt your usual blood glucose control and could result in hypoglycaemia (low blood glucose) or hyperglycaemia (high blood glucose).

Preparing for a procedure and responding to changes to your blood glucose following your procedure can reduce your risk of infections. Your diabetes team can help you during your admission and support your safe discharge home.

How can I prepare for my procedure?

Please bring with you to the hospital:

- All your medications and a current list.
- Your blood glucose/ketone meter, continuous glucose monitor (CGM), relevant consumables, your glucose diary and/or CGM system report. In most instances, you can continue to use your own equipment.
- Your injecting device/s and additional consumables. You may like to use your own insulin pen/s before or after your procedure.
- A copy of your *Hypoglycaemia Action Plan* and *Hyperglycaemia/Sick Day Action Plan*.

Continue to check your blood glucose. If you have low blood glucose, follow your *Hypoglycaemia Action Plan*. If you have high blood glucose, follow your *Hyperglycaemia/Sick Day Action Plan*.

Please ask a family member or friend to bring you. Do not drive yourself to your hospital admission.

I do not use diabetes medication. How does this affect me?

Usually there are no special requirements to follow.

Are there specific instructions for diabetes tablets and/or injectables?

Yes, an endoscopy requires a period of fasting and specific adjustments to your tablets and/or injectables on the day of the procedure.

What to do on the day prior to the procedure?

For diabetes tablets and (non-insulin) injectables

- Continue your usual oral diabetes medications (e.g. Metformin or sodium glucose co-transporter 2 (SGLT2) inhibitor).
- Continue your usual (non-insulin) injectables (e.g. dulaglutide (Trulicity®), exenatide (Bydureon®) and liraglutide (Victoza®)).

For long acting (basal) insulin

- Continue the usual insulin dose/s OR
- Reduce your evening basal insulin dose by 20% if you have had recent overnight hypoglycaemia.

For pre-mixed insulin

- Continue your usual insulin dose/s OR
- Reduce your evening insulin dose by 20% if you have had recent overnight hypoglycaemia

For co-formulation insulin

- Continue your usual insulin dose/s OR
- Reduce your evening insulin dose by 20% if you have had recent overnight hypoglycaemia

For rapid acting (mealtime) insulin

- Continue your usual mealtime bolus insulin doses.

- Continue to check your blood glucose at your usual times and if concerned.
- If you have low blood glucose, follow your *Hypoglycaemia Action Plan*.
- If you have high blood glucose, follow your *Hyperglycaemia Action Plan*.

What to do on the day of the procedure?

For diabetes tablets and (non-insulin) injectables

- Do not take your diabetes tablets.
- Do not administer your (non-insulin) injectables.

For long acting (basal) insulin

- Continue your usual insulin dose/s.

For pre-mixed insulin

- If your procedure is in the morning, you will be fasting from 6:00am: reduce your usual breakfast dose by 50%.
- If your procedure is in the afternoon, you will be fasting from 11:00am after a light breakfast: reduce your usual breakfast dose by 50%.

For co-formulation insulin

- If your procedure is in the morning, you will be fasting from 6:00am: do not take your usual breakfast insulin dose.
- If your surgery is in the afternoon, you will be fasting from 11:00am after a light breakfast: reduce your usual breakfast dose by 50%.

For rapid acting (mealtime) insulin

- If your procedure is in the morning, you will be fasting from 6:00am: do not take your breakfast bolus insulin as you will not be eating.
- If your procedure is in the afternoon, you will be fasting from 11:00am after a light breakfast: reduce your usual breakfast dose by 50%.

- Check your blood glucose every 2-4hours from the time you wake until the time you arrive at the hospital.
- If you have low blood glucose, follow your *Hypoglycaemia Action Plan*.
- If you have high blood glucose, follow your *Hyperglycaemia/Sick Day Action Plan* which may include blood ketone testing and correction bolus insulin dose instructions.

What will happen when I am admitted?

Please inform medical and nursing staff of any of the following:

- hypoglycaemia and treatment used
- hyperglycaemia and action taken.

The medical and nursing staff will check your blood glucose. If your blood glucose is above 10.0mmol/L, a correction bolus insulin dose may be used to return your blood glucose levels to target, aid recovery and assist your body to fight infection.

Where possible, self-care of your injectable device/s are encouraged and supported. Regional hospitals require people with type 2 diabetes using injectable device/s to use the supplied safety pen needles. Your medical and nursing staff and diabetes team need to know what injecting device/s you are using so that they can prevent complications and assist you in your recovery.

What will happen after my procedure?

Your diabetes tablets, (non-insulin) injectables and/or insulin injections will be restarted as soon as possible after your procedure. This is usually when you are comfortably eating and drinking again.

What support do I have on discharge?

The medical and nursing staff will assist you to restart your medications and plan your discharge. They will also be available after you are discharged home to monitor your recovery and discuss any concerns that you may have.

Your diabetes team are available to discuss your return to your usual diabetes management or provide alternative instructions. If required, your diabetes team can arrange a follow up appointment to review your diabetes management after discharge.

Please ask a family member or friend to take you home. Do not drive yourself.

Additional information

Where can I get more information?

- Diabetes Australia www.diabetesaustralia.com.au
- National Diabetes Services Scheme www.ndss.com.au
- Juvenile Diabetes Research Foundation www.jdrf.org.au
- My D (for under 25s) www.ndss.com.au/MyD

For more information

Rural Support Service Diabetes Service

PO Box 3017, Rundle Mall
ADELAIDE SA 5000

Email: Health.DiabetesService@sa.gov.au

www.chsa-diabetes.org.au

www.sahealth.sa.gov.au/regionalhealth

Public-13-A2

